Mon. Not. R. Astron. Soc. **000**, 1–7 (2000) Printed 1 February 2008 (MN LATEX style file v1.4)

The statistics of wide-separation lensed quasars

Daniel J. Mortlock[1,2,3⋆] and Rachel L. Webster[1⋆]

[1] *School of Physics, The University of Melbourne, Parkville, Victoria 3052, Australia*
[2] *Astrophysics Group, Cavendish Laboratory, Madingley Road, Cambridge CB3 0HE, U.K.*
[3] *Institute of Astronomy, Madingley Road, Cambridge CB3 0HA, U.K.*

arXiv:astro-ph/0008081v1 4 Aug 2000

Accepted. Received; in original form 2000 April 19

ABSTRACT

The absence of any wide-separation gravitational lenses in the Large Bright Quasar Survey is used to place limits on the population of cluster-sized halos in the universe, and hence constrain a number of cosmological parameters. The results agree with previous investigations in strongly ruling out the standard cold dark matter model but they are consistent with low-density universes in which the primordial fluctuation spectrum matches both cluster abundances and cosmic microwave background measurements. These conclusions are essentially independent of the cosmological constant, which is in stark contrast to the statistics of galaxy lenses. The constraints presented here are nullified if clusters have core radii of $\gtrsim 10$ kpc, but are free of a number of potential systematic errors, due to the homogeneity of the data.

Key words: gravitational lensing – cosmology: theory – galaxies: clusters.

1 INTRODUCTION

The fraction of quasars which are gravitationally-lensed by intervening mass concentrations is a function of the deflector population and the underlying cosmological model, which can then be constrained from the observed lensing frequency (e.g. Kochanek 1993, 1996). In most cases of quasar lensing known to date the principal deflectors are elliptical galaxies (e.g. Keeton, Kochanek & Seljak 1997), and the number of such lenses, together with the distribution of image separations and deflector redshifts, has been used to place quite stringent limits on the galaxy population (e.g. Kochanek 1993) and the cosmological model (e.g. Kochanek 1996; Falco, Kochanek & Muñoz 1998). Unfortunately, the dependencies of the expected lensing probability on the galaxy population and the cosmology cannot be easily separated. For instance, cosmological constant-dominated models overpredict the number of lenses if the local and cosmological galaxy populations are similar, but can be made consistent with observations if high-redshift galaxies are optically thick due to dust (e.g. Kochanek 1996; Malhotra, Rhoads & Turner 1997).

One way to bypass the lack of knowledge about the high-redshift galaxy population is to concentrate on lenses with larger image separations (i.e. $\gtrsim 3$ arcsec). These can only be produced by groups and clusters of galaxies, a population which is more closely linked to the underlying cosmological model. Just four such wide-separation lenses have been confirmed: Q 0957+561 (Walsh, Carswell & Weymann 1979); MG 2016+112 (Lawrence et al. 1984); HE 1104−1805 (Wisotzki et al. 1993); and RXJ 0911+0551 (Bade et al. 1997). There are more than ten other candidates, but statistical arguments suggest that almost all of them are physically distinct binary quasars (Kochanek, Falco & Muñoz 1999; Mortlock, Webster & Francis 1999). If this interpretation is correct, less than 0.1 per cent of all quasars are lensed with such large image separations. Conversely, theoretical calculations based on N-body simulations (e.g. Wambsganss et al. 1995) and analytical models (e.g. Narayan & White 1988; Kochanek 1995; Mortlock, Hewett & Webster 1996) predict more wide-separation lenses than are observed, if a standard cold dark matter (CDM) cosmology is assumed. However models with either a non-zero cosmological constant or strong biasing (allowing the underlying dark matter distribution to be much smoother than the galaxy distribution) are consistent with both lensing results and cosmic microwave background (CMB) anisotropies (Kochanek 1995).

The main limitations on these results are the lack of knowledge of cluster core radii (If they are greater than ~ 10 kpc, the standard models can be reconciled with the data.) and the heterogeneous nature of quasar catalogues used in previous analyses. Both these points are addressed here, with the use of a homogeneous lens survey (Section 2), and the possibility of a finite core included explicitly in the lens model (Section 3). The absence of any lenses in the data can then be used to directly constrain a number of cosmological parameters (Section 4). The accuracy of these inferences

⋆ E-mail: mortlock@ast.cam.ac.uk (DJM); rwebster@physics.unimelb.edu.au (RLW)

© 2000 RAS

is limited by both the simple model used for the deflector population and the size of the quasar sample, as discussed in Section 5.

2 THE LARGE BRIGHT QUASAR SURVEY

The Large Bright Quasar Survey (LBQS; Hewett, Foltz & Chaffee 1995) is a sample of 1055 quasars taken from 18 United Kingdom Schmidt Telescope (UKST) fields. The resultant magnitude-limited object catalogues were then used in combination with objective-prism plates to generate a list of quasar candidates. As such, the quasar catalogue was generated without any explicit morphological selection. The magnitude limit[†] varies from plate to plate, ranging from $m_{\rm lim} = 18.41$ to $m_{\rm lim} = 18.85$, and there is also a low redshift cut-off at $z = 0.2$.

The specific form of the differential number counts used in the calculation of the lensing magnification bias (See Section 3.3.) is

$$\frac{{\rm d}^2 N_{\rm q}}{{\rm d}m{\rm d}z} \propto \frac{1}{10^{\alpha_{\rm q}(m-m_{\rm q0})} - 10^{\beta_{\rm q}(m-m_{\rm q0})}}, \qquad (1)$$

where $m_0 = 19.0 \pm 0.2$ is the quasar break magnitude, $\alpha_{\rm q} = 0.9 \pm 0.1$ the bright-end slope and $\beta_{\rm q} = 0.3 \pm 0.1$ the faint-end slope. This functional form is taken from Boyle, Shanks & Peterson (1988) and Kochanek (1996), but the parameter values are chosen to match the LBQS (for $m \lesssim 19$) and the faint-end slope of the compilation of data presented by Hartwick & Schade (1990). Whilst Hewett, Foltz & Chaffee (1993) found some discrepancies between the LBQS counts and the Boyle et al. (1988) sample, the simple parameterisation of equation (1) is sufficiently accurate in the context of this calculation.

As part of the LBQS there was a systematic search for companions within ~ 10 arcsec of each quasar. The quality of the data used – the same UKST plates – is such that most companions with $m_{B_{\rm J}} \lesssim 21.5$ are found (Hewett et al. 1998), but the search was incomplete for image separations $\Delta\theta \lesssim 3$ arcsec due to the point spread function of the plates. Given the LBQS magnitude limit of ~ 18.5, the companion search has sufficient dynamic range to easily pick out most secondary lensed images, and so can be considered complete in the annulus between 3 arcsec and 10 arcsec. It is possible for galactic lenses to produce image separations in this range, but only if they are both nearby and very massive, and thus easily detectable. Hence the LBQS can be used to unambiguously constrain the population of galaxy group and cluster lenses.

The search for neighbouring images has yielded five quasar pairs so far, and it is 'unlikely that further pairs will be identified' (Hewett et al. 1998). Two of these pairs have vastly different redshifts, and are not the result of lensing (although their existence is evidence of the effectiveness of the companion search); the other three are either lenses or physical binary quasars. Q 1009−0252 (Hewett et al. 1994) is a gravitational lens, but, with $\Delta\theta = 1.5$ arcsec, it cannot be included in the lens calculation as the small-separation lens search is not well characterised. The second pair, Q 1429−0053 (Hewett et al. 1989) is a potential wide-separation lens, with an image separation of 5.1 arcsec and apparently similar spectra. However principal components analysis (e.g. Murtagh & Hecht 1987) shows that the two spectra are no more alike than a pair of spectra chosen at random from the survey (Mortlock et al. 1999). Combined with the absence of any potential deflector and the statistical arguments of Kochanek et al. (1999), it is highly unlikely that Q 1429−0053 is a lens. The third pair, Q 2153−2056 (Hewett et al. 1998), has an image separation of 7.8 arcsec, but very different spectra, and is even less likely to be a lens. Thus the LBQS almost certainly represents a sample of over 1000 quasars that is devoid of wide-separation lenses.

3 THE CALCULATION

A given world model can be characterised by one number: P_0, the probability that no wide-separation lenses are observed in the LBQS. This, in turn, is simply the product of the probabilities that each individual quasar is unlensed, so that

$$P_0 = \prod_{q=1}^{N_{\rm q}} (1 - p_q) \simeq (1 - \langle p_{\rm q} \rangle)^{N_{\rm q}}, \qquad (2)$$

where $N_{\rm q}$ is the number of quasars in the survey, p_q is the probability that the qth quasar is lensed, and $\langle p_{\rm q} \rangle$ is the lensing probability averaged over the survey. Note that this is a cumulative probability, and so the normalisation is unambiguous.

The likelihood of a given model is identified with P_0, thus ignoring any prior information. Hence, from equation (2), the only models that can be rejected at the 99 per cent level by the absence of lenses in the LBQS are those for which $\langle p_{\rm q} \rangle \gtrsim 0.004$. This is true of a number of popular cosmological scenarios, and so the data at hand are far from redundant in this context.

Having set out the statistical framework for the calculation, the next step is to define the populations and mass distributions (Sections 3.1 and 3.2, respectively), from which the lensing probability can be calculated (Section 3.3).

3.1 Deflector population

The population of collapsed halos can most accurately be estimated from N-body simulations (e.g. White et al. 1987; Efstathiou et al. 1988), but this approach is too computationally expensive to explore a wide range of cosmological models[‡]. The obvious alternative is the analytical Press-Schechter (1974) formalism (e.g. Peacock 1999). The halo population is normally given as a co-moving mass function, but the isothermal sphere lens model used in Section 3.2 is parameterised by its line-of-sight velocity dispersion, σ,

[†] All magnitudes are in the $B_{\rm J}$ system, but this subscript is omitted for brevity.

[‡] The cosmological model is defined by $\Omega_{\rm m_0}$, the present day normalised density of the universe, Ω_{Λ_0}, the current value of the similarly-normalised cosmological constant, and Hubble's constant, $H_0 \simeq 70$ km s^{-1} Mpc^{-1}.

rather than mass. Equating the total mass to that inside the virial radius of the isothermal sphere (e.g. Narayan & White 1988; Kochanek 1995) results in the conversion

$$M(\sigma,z)=\frac{\left(2^{1/2}\sigma\right)^3}{10GH(z)}, \quad (3)$$

where G is Newton's constant and $H(z)=H_0[\Omega_{\rm m_0}(1+z)^3+\Omega_{\Lambda_0}-(\Omega_{\rm m_0}+\Omega_{\Lambda_0}-1)(1+z)^2]^{1/2}$. The final expression for the co-moving halo population is then (Mortlock 1999)

$$\begin{aligned}\frac{{\rm d}n_{\rm d}}{{\rm d}\sigma}(z) &= -\frac{15\Omega_{\rm m_0}H_0^3}{8\pi^{3/2}}\frac{H(z)/H_0\,\delta_{\rm crit}(z)}{\sigma^4\,\Delta\left[M(\sigma,z)\right]}\\ &\times \frac{{\rm d}\ln\left\{\Delta\left[M(\sigma,z)\right]\right\}}{{\rm d}\ln(\sigma)}\exp\left\{\frac{\delta_{\rm crit}^2(z)}{2\Delta^2\left[M(\sigma,z)\right]}\right\}, \quad (4)\end{aligned}$$

where $M(\sigma,z)$ is given in equation (3). The cosmological model enters equation (4) only through $\delta_{\rm crit}(z)$, the extrapolated linear overdensity that would have collapsed at redshift z (e.g. Peebles 1980). The present day variance on mass scale M, $\Delta(M)$, is determined by the power spectrum of density fluctuations. The approximate CDM power spectrum of Efstathiou, Bond & White (1992) is adopted here. Most of the power spectrum parameters have little influence on the lensing likelihood (Kochanek 1995; Mortlock 1999), and so only the primordial power-law slope, n, and the normalisation, Δ_8 (the present day variance in spheres of radius 8 Mpc), are allowed to vary.

3.2 Lens model

The mass distributions of galaxy clusters – the inner regions in particular – are not well constrained by observations or theory: CDM-based N-body simulations suggest that they are singular (e.g. Navarro, Frenk & White 1997); the properties of giant arcs imply they have small core radii (e.g. Fort et al. 1992; Smail et al. 1995, but see also Bartelmann 1996) or very massive central galaxies (Williams, Navarro & Bartelmann 1999); and other observational data span the possibilities (e.g. Narayan, Blandford & Nityananda 1984; Mohr et al. 1996; Pointecouteau et al. 1999). A simple isothermal sphere (e.g. Turner, Ostriker & Gott 1984; Binney & Tremaine 1987) is adopted here because the principal analytic alternative – the Navarro et al. (1997) profile – is inconsistent with the strong lensing properties of clusters (Williams et al. 1999). However the main short-coming of both these models is the absence of any substructure, the inclusion of which increases image separations by ~ 50 per cent on average (Bartelmann, Steinmetz & Weiss 1995).

The isothermal profile, parameterised primarily its velocity dispersion, is assumed to have a core radius, $r_{\rm c}$, which scales as $r_{\rm c}=r_{\rm c*}(\sigma/\sigma_*)^{u_{\rm c}}$, where $\sigma_*=1000$ km s^{-1} is chosen arbitrarily. Its lensing properties are usually cast in terms of the singular model's Einstein radius, $\theta_{\rm E}$ (which is dependent on the cosmological model; Schneider, Ehlers & Falco 1992), and the critical radius in the lens plane, $\beta_{\rm crit}$ (which decreases as the core radius increases). For the definitions of these terms and a further discussion of this lens model see, Hinshaw & Krauss (1984), Kochanek (1996) or Mortlock & Webster (2000).

3.3 Lensing probability

From the properties of the lens model it is possible to calculate the probability, $p_{\rm q}$, that a quasar of magnitude $m_{\rm q}$ (in the parent survey) and redshift $z_{\rm q}$ is found to be lensed. This is derived rigorously for a typical lens survey in Kochanek (1996) and Mortlock & Webster (2000), but there are some important differences in the calculation of the magnification bias here. In most lens surveys, the angular separation of the images is less than the angular resolution of the parent survey, and so, for a lensed quasar, $m_{\rm q}$ is given by summing the fluxes of all the images. However, in the case of the UKST data from which the LBQS was selected, only images separated by $\lesssim 5$ arcsec appear merged (Webster, Hewett & Irwin 1988, although Hewett et al. 1995 give ~ 6 arcsec; this discrepancy leads to a ~ 10 per cent uncertainty in the calculated lensing probability), whereas the search annulus of the lens survey extends from $\Delta\theta_{\rm min}=3$ arcsec to $\Delta\theta_{\rm max}=10$ arcsec. This is an unusual situation – the lens survey improves on the parent survey in depth, not resolution. For any lenses with $\Delta\theta\lesssim 5$ arcsec the standard magnification bias is correct, but for wider-separation multiples $m_{\rm q}$ is determined only by the magnification of the brightest individual image[§]. This is often considerably less than the total magnification, and the resultant reduction in the lensing probability can be quite marked, as shown below.

The first step in evaluating $p_{\rm q}$ is to calculate $p_{\rm q,d}$, the probability that a given quasar is lensed by a particular halo, defined by its redshift, $z_{\rm d}$ and its velocity dispersion, σ. With the normal total magnification bias, this is given by

$$p_{\rm q,d}=\frac{\int_0^{\beta_{\rm crit}}2\pi\beta\,S(\beta)\left.\frac{{\rm d}^2N_{\rm q}}{{\rm d}z_{\rm q}\,{\rm d}m}\right|_{m=m_{\rm q}+5/2\,\log[\mu_{\rm tot}(\beta)]}{\rm d}\beta}{4\pi\left.\frac{{\rm d}^2N_{\rm q}}{{\rm d}z_{\rm q}\,{\rm d}m_{\rm q}}\right|}, \quad (5)$$

where the quasar luminosity function, ${\rm d}^2N_{\rm q}/{\rm d}z\,{\rm d}m$, is given in equation (1) and $S(\beta)$ is the selection function (c.f. Kochanek 1996). The latter is approximated by

$$\begin{aligned}S(\beta) &= H[\Delta m_{\rm max}-\Delta m(\beta)]\\ &\times H[\Delta\theta(\beta)-\Delta\theta_{\rm min}]\,H[\Delta\theta_{\rm max}-\Delta\theta(\beta)]\,, \quad (6)\end{aligned}$$

where $H(x)$ is the Heavyside step function.

Equation (5) can be converted to the single image magnification bias simply by replacing $\mu_{\rm tot}(\beta)$ with the magnification of the brightest image, $\mu_{\rm max}(\beta)$. As $\mu_{\rm max}<\mu_{\rm tot}$ in all cases of multiple imaging, the single image magnification bias always results in a lower value of $p_{\rm q,d}$. The ratio of the two probabilities is particularly simple to calculate if the deflectors are assumed to be singular: the integration variable in equation (5) can be changed to $\mu_{\rm tot}=2/(\beta/\theta_{\rm E})$, and $\mu_{\rm max}=(1+\beta/\theta_{\rm E})/(\beta/\theta_{\rm E})$. Thus

$$\frac{p_{\rm q,d\,single}}{p_{\rm q,d\,total}}=\frac{\int_2^\infty\frac{2}{(\mu_{\rm max}-1)^3}\left.\frac{{\rm d}^2N_{\rm q}}{{\rm d}z_{\rm q}\,{\rm d}m}\right|_{m=m_{\rm q}+5/2\,\log(\mu_{\rm max})}{\rm d}\mu_{\rm max}}{\int_2^\infty\frac{8}{\mu_{\rm tot}^3}\left.\frac{{\rm d}^2N_{\rm q}}{{\rm d}z_{\rm q}\,{\rm d}m}\right|_{m=m_{\rm q}+5/2\,\log(\mu_{\rm tot})}{\rm d}\mu_{\rm tot}}, \quad (7)$$

where $\Delta m_{\rm max}\to\infty$ for simplicity. Fig. 1 shows how this ratio varies with the slope of the quasar number counts,

[§] Merging images are not explicitly treated here.



Figure 1. The effect of the single image magnification bias (relative to the standard total magnification bias) on the lensing probability of a single deflector, $p_{\rm q,d}$, as a function of the logarithmic slope of the source counts, $\alpha_{\rm q}$. The bright end and faint end slopes inferred from the Boyle et al. (1988) quasar sample are also shown. The deflector (which has a velocity dispersion of $\sigma = 1000$ km s^{-1}) and source are at redshifts of 0.5 and 2, respectively, and $\Omega_{\rm m_0} = 1$ and $\Omega_{\Lambda_0} = 0$ is assumed. Results are shown for several core radii: $r_{\rm c} = 0$ kpc (singular model; solid line); $r_{\rm c} = 5$ kpc (dashed line); $r_{\rm c} = 10$ kpc (dot-dashed line) and $r_{\rm c} = 20$ kpc (dotted line).

$\alpha_{\rm q}$, if they are given by a single power law, $\mathrm{d}N_{\rm q}/\mathrm{d}m \propto 10^{\alpha_{\rm q} m}$. The main difference between the two situations is the reduction in the amplitude of the high-magnification tail in the case of the single image magnification bias, which in turn has a greater effect on $p_{\rm q,d}$ if the luminosity function is steeper. Thus the wide-separation lensing probability for bright quasars (including the LBQS) is up to a factor of three lower than previous calculations would suggest¶.

The probability that a quasar is lensed by any halo is obtained by integrating $p_{\rm q,d}$ over the deflector population, to give

$$p_{\rm q} = \int_0^{z_{\rm q}} \int_0^{\infty} \left.\frac{\mathrm{d}V_0}{\mathrm{d}z}\right|_{z=z_{\rm d}} \frac{\mathrm{d}n_{\rm d}}{\mathrm{d}\sigma}(z)\, p_{\rm q,d}\, \mathrm{d}\sigma\, \mathrm{d}z_{\rm d}, \qquad (8)$$

where $\mathrm{d}n_{\rm d}/\mathrm{d}\sigma(z)$, the co-moving density of deflectors, is given in Section 3.1.

This must be calculated separately for each quasar in the LBQS, due mainly to their large redshift range, and the resultant likelihoods can be averaged to give $\langle p_{\rm q} \rangle$. This is shown in Figs. 2 and 3 as function of several model parameters. In Fig. 2 (a), the lensing probability initially increases very rapidly with Δ_8, as the velocity dispersion of the largest halos depends exponentially on Δ_8 (Kochanek 1995). For $\Delta_8 \gtrsim \delta_{\rm crit}$, however, the lensing likelihood flattens off as the largest halos produce image separations of $\gtrsim 10$ arcsec,

¶ Most relevantly, Kochanek (1995) used the normal magnification bias, but due to the image separations under consideration (up to 1 arcmin), the weaker, single image bias would have given a more realistic value of the lensing probability. For example, the two standard deviation result that $0.27 \lesssim \Delta_8 \lesssim 0.63$ (in the standard CDM model) becomes $0.35 \lesssim \Delta_8 \lesssim 0.75$.



Figure 2. The average probability that any quasar in the LBQS is observed to be lensed, $\langle p_{\rm q} \rangle$. The dependence on the normalisation, Δ_8, is shown in (a); the variation with the canonical core radius, $r_{\rm c*}$ is shown in (b); and the dependence on the slope of the power spectrum, n, is shown in (c). The 'default' values are $\Delta_8 = 1$, $r_{\rm c*} = 0$ and $n = 1$ and the three lines represent different cosmological models: $\Omega_{\rm m_0} = 1.0$ and $\Omega_{\Lambda_0} = 0.0$ (solid lines); $\Omega_{\rm m_0} = 0.3$ and $\Omega_{\Lambda_0} = 0.0$ (dashed lines); and $\Omega_{\rm m_0} = 0.3$ and $\Omega_{\Lambda_0} = 0.7$ (dot-dashed lines). Models with $\langle p_{\rm q} \rangle > 0.004$ (the horizontal dashed line) can be rejected at the 99 per cent level.

and so cannot contribute to $\langle p_{\rm q}\rangle$. As shown in Fig. 2 (b), the number of lenses decreases with cluster core radii, as expected (e.g. Kochanek 1995). However the dependence is very weak if $u_{\rm c} \simeq 4$. For a given $r_{\rm c*}$, the smaller deflectors (which are capable of producing lenses with $\Delta\theta \simeq 3$ arcsec) are more effective, being nearly singular. Unfortunately this degeneracy means that any limits placed on the other model parameters are weakened by the fact that $u_{\rm c}$ is essentially unconstrained. The situation with respect to n is more promising – as can be seen from Fig. 2 (c) the slope of the power spectrum has only a small effect on the lens statistics.

The dependence of $\langle p_{\rm q}\rangle$ on the cosmological model is illustrated in both Fig. 2 and Fig. 3. The most striking aspect of these plots is that $\Omega_{\rm m_0}$ is considerably more important than Ω_{Λ_0}, whereas it is the cosmological constant that dominates the statistics of lensing by galaxy-scale objects. For a given $\Omega_{\rm m_0}$ the volume element is greater in high-Ω_{Λ_0} universes, but clusters form at later times, so the increase of $\langle p_{\rm q}\rangle$ with Ω_{Λ_0} is only mild. The strong dependence on $\Omega_{\rm m_0}$ comes about primarily as halos are heavier for a fixed Δ_8. A cluster that has collapsed from an initial perturbation of a given co-moving scale has mass $M \propto \Omega_{\rm m_0}$, velocity dispersion $\sigma \propto \Omega_{\rm m_0}^{1/3}$ [from equation (3)], and hence a lensing cross-section that increases as $\Omega_{\rm m_0}^{4/3}$ (e.g. Turner et al. 1984; Kochanek 1995). This dependence is apparent for low-density models (See Fig. 3.), but as $\Omega_{\rm m_0} \to 1$ the predominant halos have typical image separations that are greater than $\Delta\theta_{\rm max}$. A lens survey with a broader search annulus than that of the LBQS would be required to probe models with either Δ_8 or $\Omega_{\rm m_0}$ of order unity.



Figure 3. The probability that any LBQS quasar is lensed, $\langle p_{\rm q}\rangle$, as a function of $\Omega_{\rm m_0}$ and Ω_{Λ_0}. The power spectrum is parameterised by $\Delta_8 = 1$, and $n = 1$, and the lenses are singular. The contours start at $\langle p_{\rm q}\rangle = 0$ (the $\Omega_{\rm m_0} = 0$ axis) and then increase in steps of 0.005; the highest contour (at the top-right) is $\langle p_{\rm q}\rangle = 0.03$. The dashed line denotes spatially flat models.

4 RESULTS

The selection criteria for a lens to appear in the LBQS (Section 2) and the lensing calculation described in Section 3 can now be combined to give P_0, the probability that the LBQS contains no wide-separation lenses. This is shown for several combinations of model parameters in Fig. 4. The contours shown are for $P_0 = 0.01$ (i.e. 99 per cent limits), $P_0 = 0.05$ and $P_0 = 0.5$, but are only one-sided as models which predict an arbitrarily low number of lenses are perfectly consistent with the data.

Fig. 4 (a) shows that $\Delta_8 \lesssim 0.4$ (with 99 per cent confidence) in an Einstein-de Sitter (EdS) cosmology with a standard CDM spectrum. This is considerably lower than the value of 1.4 ± 0.1 inferred from the *Cosmic Background Explorer* (*COBE*) data (Smoot et al. 1992). Standard *COBE*-normalised CDM models can be made consistent with the data if $r_{\rm c*} \gtrsim 10$ kpc and $u_{\rm c} \simeq 2$, as shown in Fig. 4 (b). Although such models are somewhat contrived[‖], the scaling of cluster core radii is so poorly constrained that they cannot be completely ruled out. However it is the same low-density models implied by observations of high-redshift supernovæ (e.g. Schmidt et al. 1998; Perlmutter et al. 1999) and the position of the first CMB Doppler peak (e.g. Efstathiou et al. 1999; de Bernardis et al. 2000) that are favoured by the data at hand. There may be a slight discrepancy if measurements of Δ_8 are also considered, as the values inferred from the local cluster population (e.g. Peebles 1989; Frenk et al. 1990; Bahcall, Fan & Cen 1997) and the galaxy-galaxy correlation function (e.g. Maddox et al. 1990; Efstathiou et al. 1992) are slightly higher than allowed in the $\Omega_{\rm m_0} = 0.3$ model here. However, if $\Omega_{\rm m_0} \simeq 0.2$ and the universe is flat then the greatest source of disagreement with the absence of wide-separation lenses in the LBQS is, somewhat ironically, the low number of small-separation lenses that preclude $\Omega_{\Lambda_0} \simeq 0.8$ (e.g. Kochanek 1996).

It is possible, although very unlikely, that there is a wide-separation lens in the LBQS, as Q 1429−0053 has not been conclusively proved to be a binary quasar and there is still a small amount of the companion search to be completed (Hewett et al. 1998). If such a lens were to be discovered then the limits on the model parameters would be shifted somewhat. Performing a calculation similar to that described in Section 3 gives the probability of finding one lens in the LBQS as

$$\begin{aligned} P_1 &\simeq N_{\rm q}\langle p_{\rm q}\rangle(1-\langle p_{\rm q}\rangle)^{N_{\rm q}-1} \\ &\simeq N_{\rm q} P_0^{(N_{\rm q}-1)/N_{\rm q}}(1-P_0)^{1/N_{\rm q}}, \end{aligned} \qquad (9)$$

where $N_{\rm q} = 1055$ is the number of quasars in the survey. From this conversion, the 1 per cent, 5 per cent and 50 per cent contours for P_0 shown in Fig. 4 become 5 per cent, 15 per cent and 35 per cent contours for P_1, respectively. In other words, the 99 per cent limits would be weakened to 95 per cent limits. The constraints would also become

[‖] There must also be some nearly-singular clusters to account for the morphology of the confirmed wide-separation lenses listed in Section 1. Observations of giant arcs (e.g. Smail et al. 1995; Bartelmann 1996) also argue against such high values of $_{\rm c*}$.



Figure 4. The constraints placed on the cosmological model and the power spectrum parameters from the absence of lenses in the LBQS. The solid contours are at $P_0 = 0.01$; the dashed contours are at $P_0 = 0.05$ and $P_0 = 0.5$. In all panels the vertical axis gives the normalisation of the power spectrum, Δ_8. The horizontal axes are: Ω_{m_0} (for both $\Omega_{\Lambda_0} = 0$ and flat models) in (a); the canonical core radius, r_{c*} (for $u_c = 2$ and $u_c = 4$) in (b); and the slope of the power spectrum, n, in (c). In panels (b) and (c) an EdS cosmology is assumed, and $r_{c*} = 0$ and $n = 1$ are assumed except where these parameters are explicitly varied.

two-sided, but the models that would be ruled out by the presence of a lens have $\langle p_q \rangle \simeq 10^{-5}$, and would not be considered plausible a priori. Thus the presence of a wide-separation lens in the LBQS would broaden the constraints on the model parameters, but would not qualitatively change the conclusions.

5 CONCLUSIONS

The fact that no wide-separation ($\Delta\theta \gtrsim 3$ arcsec) lenses have been discovered in the LBQS places tight constraints on the population of cluster-mass objects in the universe. Assuming a hierarchical theory of halo formation, one-sided limits can then be placed on several cosmological parameters. Interestingly, the results presented in Section 4 (See also Kochanek 1995.) indicate that only three model parameters are strongly constrained (Ω_{m_0}, Δ_8 and r_{c*}), and that the lens statistics are almost independent of the cosmological constant. This is in stark contrast with the probability of lensing by galaxies, which is dominated by Ω_{Λ_0}.

The most straightforward result is that standard the CDM model (i.e. an EdS cosmology with a power spectrum index of $n = 1$) is completely ruled out unless the normalisation is very low ($\Delta_8 \lesssim 0.4$ at the 99 per cent level). Only if the core radii of $\sigma \simeq 1000$ km s^{-1} clusters are $\gtrsim 10$ kpc, and $r_c \propto \sigma^2$ can these limits be avoided. For non-EdS cosmologies, the lensing results are in agreement with most a priori reasonable models (as discussed in Section 4), although $\Omega_{m_0} \lesssim 0.2$ is implied unless Δ_8 is low or, again, clusters have large core radii.

Whilst the conclusions are in agreement with most expectations, it is important to consider the various approximations in this calculation, and their effects on the parameter constraints. Fortunately, the lensing probability varies as $\Omega_{m_0}^{4/3}$ and increases exponentially with Δ_8 (provided $\Delta_8 \lesssim 1$), whereas the use of the Press-Schechter (1974) formalism, the calculation of δ_{crit}, the choice of a smooth, spherical lens model, and the treatment of the magnification bias all lead to uncertainties of tens of per cent at most. Hence the limits on Ω_{m_0} and Δ_8 could be changed by similar amounts. The choice of mass profile used for the lenses is more important (as illustrated by the effect of r_{c*}) and it would be especially interesting to perform the above calculation using the Navarro et al. (1997) halo model, as the mass distribution of the halos in their prescription is assumed to be directly related to the underlying cosmological model.

The LBQS represents an ideal type of data-set for this kind of analysis, as it has well-characterised selection effects and the companion search was both deep and systematic; the only real shortcoming is the size of the survey. Both the Jodrell Bank-Very Large Array Astrometric Survey (JVAS; Patnaik et al. 1992) of ~ 2500 sources and the smaller *Hubble Space Telescope* Snapshot Survey (Maoz et al. 1997) of ~ 500 quasars also contain no confirmed lenses with $\Delta\theta \gtrsim 5$ arcsec (Marlow et al. 1998), and so would imply similar one-sided limits to those presented here. The situation should be improved with the advent of much larger surveys, such as the 2 degree Field (2dF) quasar survey (e.g. Boyle et al. 1999a,b), with $\sim 3 \times 10^4$ quasars, and the Sloan Digital Sky Survey (SDSS; e.g. Szalay 1998; Loveday & Pier 1998), with $\sim 10^5$ quasars. The 2dF survey's companion search is

not particularly deep, and will only include lenses with image separations greater than 8 arcsec. Nonetheless, the sheer size of the data-set will allow a lensing determination of Ω_{m_0} and Δ_8 to within ~ 10 per cent. The SDSS quasar survey includes high-resolution imaging, and should thus provide a large sample of both galactic and wide-separation lenses. This will allow Ω_{m_0}, Ω_{Λ_0} and Δ_8 to be constrained simultaneously to within several per cent from lensing statistics alone.

ACKNOWLEDGMENTS

Many thanks to both Paul Hewett, whose intimate understanding of the LBQS was crucial to this analysis, and to the anonymous referee, whose comments improved the form of this paper. DJM was supported by an Australian Postgraduate Award.

REFERENCES

Bade N., Siebert J., Lopez S., Voges W., Reimers D., 1997, A&A, 317, L13
Bahcall N. A., Fan X., Cen R. Y., 1997, ApJ, 485, L53
Bartelmann M., 1996, A&A, 313, 697
Bartelmann M., Steinmetz M., Weiss A., 1995, A&A, 297, 1
Binney J. J., Tremaine S., 1987, Galactic Dynamics. Princeton University Press, Princeton
Boyle B. J., Croom S. M., Smith R. J., Shanks T., Miller L., Loaring N. S., 1999a, Phil. Trans. of the Royal Soc. A, 357, 185
Boyle B. J., Croom S. M., Smith R. J., Shanks T., Miller L., Loaring N. S., 1999b, in Morganti R., Couch W. J., eds, Looking Deep in the Southern Sky. Springer-Verlag, Berlin, p. 16
Boyle B. J., Shanks T., Peterson B. A., 1988, MNRAS, 235, 935
de Bernardis P., et al., 2000, Nature, 404, 955
Efstathiou G., Frenk C. S., White S. D. M., Davis M., 1988, MNRAS, 235, 715
Efstathiou G., Bond J. R., White, S. D. M., 1992, MNRAS, 258, 1P
Efstathiou G., Bridle S. L., Lasenby A. N., Hobson M. P., Ellis R. S., 1999, MNRAS, 303, L47
Falco E. E., Kochanek C. S., Muñoz J. A., 1998, ApJ, 494, 47
Fort B., Le Fevre O., Hammer F., Cailloux M., 1992, ApJ, 399, L125
Frenk C. S., White S. D. M., Efstathiou G., Davis M., 1990, ApJ, 351, 10
Hartwick F. D. A., Schade D., 1990, ARA&A, 28, 437
Hewett P. C., Foltz C. B., Chaffee F. H., 1993, ApJ, 406, L43
Hewett P. C., Foltz C. B., Chaffee F. H., 1995, AJ, 109, 1498
Hewett P. C., Foltz C. B., Harding M. E., Lewis G. F., 1998 AJ, 115, 383
Hewett P. C., Irwin M. J., Foltz C. B., Harding M. E., Corrigan R. T., Webster R. L., Dinshaw N., 1994, AJ, 108, 1534
Hewett P. C., Webster R. L., Harding M. E., Jedrzejewski R. I., Foltz C. B., Chaffee F. H., Irwin M. J., Le Fèvre O., 1989, ApJ, 346, L61
Hinshaw G., Krauss L. M., 1987, ApJ, 320, 468
Keeton C. R., Kochanek C. S., Seljak U., 1997, ApJ, 482, 604
Kochanek C. S., 1993, ApJ, 419, 12
Kochanek C. S., 1995, ApJ, 453, 545
Kochanek C. S., 1996, ApJ, 466, 638
Kochanek C. S., Falco E. E., Muñoz J. A., 1999, ApJ, 510, 590
Lawrence C. R., Schneider D. P., Schmidt M., Bennett C. L., Hewitt J. N., Burke B. F., Turner E. L., Gunn J. E., 1984, Science, 223, 46
Loveday J., Pier J., 1998, in Colombi S., Mellier Y., Raban B., eds, Wide Field Surveys in Cosmology. Edition Frontiers, Paris, p. 317
Maddox S. J., Efstathiou G., Sutherland W. J., Loveday J., 1990, MNRAS, 242, 43P
Malhotra S., Rhoads J. E., Turner E. L., 1997, MNRAS, 288, 138
Maoz D., Rix H.-W., Gal-Yam A., Gould A., 1997, ApJ, 486, 75
Marlow D. R., McGuinness A. D., Browne I. W. A., Wilkinson P. N., Helbig P. J., 1998, in Müller V., Gottlöber S., Mücket J. P., Wambsganss J., eds, Large Scale Structure: Tracks and Traces. World Scientific, Singapore, p. 327
Mohr J. J., Geller M. J., Fabricant D. G., Wegner G., Thorstensen J., Richstone D. O., 1996, ApJ, 470, 724
Mortlock D. J., 1999, PhD Thesis, University of Melbourne
Mortlock D. J., Webster R. L., 2000, MNRAS, in press
Mortlock D. J., Webster R. L., Francis P. J., 1999, MNRAS, 309, 836
Mortlock D. J., Webster R. L., Hewett P. C., 1996, in Kochanek C. S., Hewitt J. N., eds, Proc. IAU Symp. No. 173. Astrophysical Applications of Gravitational Lensing. Kluwer, Dordrecht, p. 71
Murtagh F., Hecht A., 1987, Multivariate Data Analysis. Reidel, Dordrecht
Narayan R., Blandford R. D., Nityananda, R., 1984, Nature, 310, 112
Narayan R., White S. D. M., 1988, MNRAS, 231, 97P
Navarro J. F., Frenk C. S., White S. D. M., 1997, ApJ, 490, 493
Patnaik A. R., Browne I. W. A., Wilkinson P. N., Wrobel J. M., 1992, MNRAS, 254, 655
Peacock J. A., 1999, Cosmological Physics. Cambridge University Press, Cambridge
Peebles P. J. E., 1980, The Large-scale Structure of the Universe. Princeton University Press, Princeton
Peebles P. J. E., 1989, Physica D, 38, 273
Perlmutter S., et al., 1999, ApJ, 517, 565
Pointecouteau E., Giard M., Benoit A., Désert F. X., Aghanim N., Coron N., Lamarre J. M., Delabrouille J., 1999, ApJ, 519, L115
Press W. H., Schechter P., 1974, ApJ, 187, 425
Schmidt B. P., et al., 1998, ApJ, 507, 46
Schneider P., Ehlers J., Falco E. E., 1992, Gravitational Lenses. Springer-Verlag, Berlin
Smail I., Couch W. J., Ellis R. S., Sharples R. M., 1995, ApJ, 440, 501
Smoot G. F., et al., 1992, ApJ, 396, L1
Szalay A. S., 1998, in Müller V., Gottlöber S., Mücket J. P., Wambsganss J., eds, Large Scale Structure: Tracks and Traces. World Scientific, Singapore, p. 97
Turner E. L., Ostriker J. P., Gott J. R., 1984, ApJ, 284, 1
Walsh D., Carswell R. F., Weymann R. J., 1979, Nature, 279, 381
Wambsganss J., Cen R., Ostriker J. P., Turner E. L., 1995, Science, 268, 274
Webster R. L., Hewett P. C., Irwin M. J., 1988, AJ, 95, 19
White S. D. M., Frenk C. S., Davis M., Efstathiou G., 1987, AJ, 313, 505
Williams L. L. R., Navarro J. F., Bartelmann M., 1999, ApJ, 527, 535
Wisotzki L., Kohler T., Kayser R., Reimers D., 1993, A&A, 278, L15

This paper has been produced using the Royal Astronomical Society/Blackwell Science LaTeX style file.